

Procter & Gamble Co.

-PE Product Labels –

-continued

DayQuil
HOT REMEDY

RAPID RELIEF
MULTI-SYMPTOM

COLD & FLU
+ CONGESTION

Acetaminophen,
Dextromethorphan HBr,
Phenylephrine HCl

NyQuil
HOT REMEDY

RAPID RELIEF
MULTI-SYMPTOM

COLD & FLU
+ CONGESTION

Acetaminophen,
Diphenhydramine HCl,
Phenylephrine HCl

HONEY LEMON
FLAVORED

Non-Drowsy

Max Strength

Nighttime Relief

10 DAYQUIL POWDER PACKETS; 6 NYQUIL POWDER PACKETS; 16 TOTAL POWDER PACKETS

how2recycle.info

MAINTAIN THE BOX

DayQuil (continued)

Other information

• acetaminophen contains propylparaben 0.1 mg per dose • do not exceed 25°C

Active ingredients (in each Packet):

Acetaminophen 10 mg • Dextromethorphan HBr 1 mg • Phenylephrine HCl 5 mg

Directions

• 10-15 years & older • 1 packet (dose) every 4 hours

Warnings

• liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Alergy Note: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Severe throat warning: If your throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, were one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, phobic or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease • heart disease
- high blood pressure • thyroid disease • diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed • do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz of hot water and stir briefly or white hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briefly before and after heating. Do not overheat.

Age

Adults & Children 12 yrs & older • 1 packet (dose) every 4 hours

Children under 12 yrs • do not use

NyQuil (continued)

Other information

• acetaminophen contains propylparaben 0.1 mg per dose • do not exceed 25°C

Active ingredients (in each Packet):

Acetaminophen 10 mg • Diphenhydramine HCl 25 mg • Phenylephrine HCl 5 mg

Directions

• 10-15 years & older • 1 packet (dose) every 4 hours

Warnings

• liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Alergy Note: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Severe throat warning: If your throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, phobic or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease • heart disease
- high blood pressure • thyroid disease • diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed • do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz of hot water and stir briefly, to white hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briefly before and after heating. Do not overheat.

Age

Adults & Children 12 yrs & older • 1 packet (dose) every 4 hours

Children under 12 yrs • do not use

DayQuil (continued)

Other information

• acetaminophen contains propylparaben 0.1 mg per dose • do not exceed 25°C

Active ingredients (in each Packet):

Acetaminophen 10 mg • Dextromethorphan HBr 1 mg • Phenylephrine HCl 5 mg

Directions

• 10-15 years & older • 1 packet (dose) every 4 hours

Warnings

• liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Alergy Note: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Severe throat warning: If your throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, were one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, phobic or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease • heart disease
- high blood pressure • thyroid disease • diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed • do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz of hot water and stir briefly, to white hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briefly before and after heating. Do not overheat.

Age

Adults & Children 12 yrs & older • 1 packet (dose) every 4 hours

Children under 12 yrs • do not use

NyQuil (continued)

Other information

• acetaminophen contains propylparaben 0.1 mg per dose • do not exceed 25°C

Active ingredients (in each Packet):

Acetaminophen 10 mg • Diphenhydramine HCl 25 mg • Phenylephrine HCl 5 mg

Directions

• 10-15 years & older • 1 packet (dose) every 4 hours

Warnings

• liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Alergy Note: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Severe throat warning: If your throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, phobic or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI,



3 23900 04198

DayQuil Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Ingredient	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Directions

Temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- runny nose & sneezing
- stuffy nose & sinus

Warnings

Lower warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- if you are now taking a prescription medicine or other drug (MDR) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 7 weeks after stopping the MDR drug. If you do not know if your prescription drug contains an MDR, ask a doctor or pharmacist before taking this product.

Other information

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Other information

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

NyQuil Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Ingredient	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Directions

Temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- runny nose & sneezing
- stuffy nose & sinus

Warnings

Lower warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- if you are now taking a prescription medicine or other drug (MDR) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 7 weeks after stopping the MDR drug. If you do not know if your prescription drug contains an MDR, ask a doctor or pharmacist before taking this product.

Other information

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Other information

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

Keep out of reach of children. See important information about acetaminophen on page 2 of this booklet.

DayQuil

ULTRA CONCENTRATED

COLD & FLU

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains, Nasal Congestion, Sinus Pressure, Cough

48 DAYQUIL LIQUICAPS™

25% SMALLER

EASY TO SWALLOW

Parental: www.dayquil.com

NyQuil

ULTRA CONCENTRATED

COLD & FLU

Acetaminophen, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains, Sneezing, Runny Nose, Cough

48 NYQUIL LIQUICAPS™ 96 TOTAL LIQUICAPS™

25% SMALLER

EASY TO SWALLOW


Parental: www.nyquil.com

Drug Facts		Drug Facts	
NyQuil™ SEVERE Cold & Flu		DayQuil™ SEVERE Cold & Flu	
Active ingredients (in each LiquiCap)	Purpose	Active ingredients (in each LiquiCap)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer	Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant	Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Sleeping aid	Doxylamine succinate 6.25 mg	Sleeping aid
Phenylephrine HCl 5 mg	Nasal decongestant	Phenylephrine HCl 5 mg	Nasal decongestant
Uses temporarily relieves common cold/flu symptoms:		Uses temporarily relieves common cold/flu symptoms:	
• nasal congestion • sinus congestion & pressure		• nasal congestion • sinus congestion & pressure	
• cough due to minor throat & bronchial irritation		• cough due to minor throat & bronchial irritation	
• cough to help you sleep		• minor aches & pains • headache • fever • sore throat	
• temporarily restores free breathing through the nose		• temporarily restores free breathing through the nose	
• reduces swelling of nasal passages		• promotes nasal and/or sinus drainage	
• temporarily restores free breathing through the nose		• helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial	
• promotes nasal and/or sinus drainage		passageways of bothersome mucus and make coughs more productive.	
Warnings		Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:		Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
• more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product		• more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product	
• 3 or more alcoholic drinks every day while using this product		• 3 or more alcoholic drinks every day while using this product	
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:		Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
• Skin redness • Blisters • Rash		• Skin redness • Blisters • Rash	
If a skin reaction occurs, stop use and seek medical help right away		If a skin reaction occurs, stop use and seek medical help right away	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	

Drug Facts (continued)		Drug Facts (continued)	
Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.		Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.	
• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.		• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland		Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin		Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin	
When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness		When using this product • do not use more than directed • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.		Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	

Drug Facts (continued)		Drug Facts (continued)	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions • take only as directed • do not exceed 8 LiquiCaps per 24 hrs		Directions • take only as directed • do not exceed 8 LiquiCaps per 24 hrs	
adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs		adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs	
children 4 to under 12 yrs ask a doctor		children 4 to under 12 yrs ask a doctor	
children under 4 yrs do not use		children under 4 yrs do not use	
Other information • do not exceed 25°C		Other information • do not exceed 25°C	
Inactive ingredients FD&C Blue No. 1, gelatin, glycerin, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution. *May contain this ingredient.		Inactive ingredients ammonium hydroxide, FD&C Blue No. 1, FD&C Red No. 40, gelatin, glycerine, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol, sorbitan solution, titanium dioxide	
Questions? 1-800-362-1663		Questions? 1-800-362-1663	





Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs
- adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs
- children 4 to under 12 yrs ask a doctor
- children under 4 yrs do not use

Other information store at no greater than 25°C

Inactive Ingredients FD&C Yellow No. 5, FD&C Yellow No. 6, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

Questions? 1-800-362-1663



DayQuil

EASY TO OPEN BOTTLE¹

ACTUAL SIZE

25% SMALLER*

*COMPARED TO ORIGINAL DAYQUIL™ 24hr BUSTER PACK

1 COMPARED TO ORIGINAL DAYQUIL™ 24hr BUSTER PACK

Drug Facts

Active ingredients (in each LiquiCap)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- itch

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

DIST. BY:
PROCTER & GAMBLE
CINCINNATI, OH 45202

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

P&G
www.pg.com
vicks.com
Puffs®
www.pg.com/patents

Headache, Fever,
Sore Throat,
Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

25% SMALLER*

EASY TO SWALLOW

48 LIQUICAPS™

3 2900 03112 **8**

90491463

MADE IN U.S.A.

PAPER BOX PLASTIC BOTTLE

Drug Facts

Active Ingredients (In each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- * temporarily relieves common cold's symptoms:
- * sinus congestion & pressure
- * minor aches & pains
- * reduces swelling of nasal passages
- * temporarily restores freer breathing through the nose
- * promotes nasal and/or sinus drainage
- * helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.
- * relief of fever
- * sore throat
- * headache
- * fever
- * chest congestion

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take
 • more than 8 caplets in 24 hours, which is the maximum daily amount for this product.
 • with other drugs containing acetaminophen. * 3 or more alcoholic drinks every day while taking this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• If you are now taking a prescription medicine like valproic acid inhibitor (MAOI) certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland

cough that occurs with too much phlegm (mucus)
 persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts

Active ingredients (in each Packet)	Purpose
Acetaminophen 500 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- minor aches & pains • sore throat • fever • headache
- cough due to minor throat & bronchial irritation • cough to help you sleep • nasal congestion
- sinus congestion & pressure • reduces swelling of nasal passages
- temporarily restores free breathing through the nose • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening • Blisters • Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days • redness or swelling is present
- new symptoms occur • cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Age	Dose
adults & children 12 yrs & over	one packet every 4 hours
children under 12 yrs	do not use

• take only as directed • do not exceed 6 doses per 24 hrs

• Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes

• If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat.

Other information • each packet contains: potassium 10 mg • phenylephrine HCl 10 mg per dose • do not exceed 25°C.

Inactive ingredients: acetylsalicylic acid, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions? 1-800-362-1683

how2recycle.info



Tamper Evident:
Do not use if inner sealed packet is torn or broken.

MADE IN MEXICO
DIST. BY:
PROCTER & GAMBLE,
CINCINNATI, OH 45202

P&G
www.pg.com

Patents: www.pg.com/patents

Keep outer package for complete product information.

91361958



50%
MORE



50%
MORE



FluTherapy^{DAY}

SEVERE COLD & FLU

Acetaminophen - Pain reliever/Fever reducer,
Dextromethorphan HBr - Cough suppressant
Phenylephrine HCl - Nasal decongestant

• Fast Relief of Cold & Flu symptoms
• Soothing Vicks Vapoars

69
PACKETS



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ACTUAL SIZE



SIDE VIEW OF 9 PACKETS

Drug Facts

Active ingredients (in each Packet)	Purpose
Acetaminophen 650 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- sore throat
- fever
- headache
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- minor aches & pains
- cough due to minor throat & bronchial irritation
- nasal congestion
- sinus congestion & pressure
- temporarily restores freer breathing through the nose
- runny nose
- sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

• with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- Do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- excitability may occur especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs

Age	Dose
Adults & children 12 yrs & over	one packet every 4 hours
children under 12 yrs	do not use

• Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes

• If using a microwave, add contents of one packet to 8 oz. of cold water, stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg
- phenylephrine: contains phenylephrine 68 mg per dose
- do not exceed 25°C.

Inactive ingredients: aceulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, tribasic calcium phosphate

Questions? 1-800-362-1683

Keep outer package for complete product information.

91380172



how2recycle.info



50% MORE



50% MORE



NIGHT FluTherapy

SEVERE COLD & FLU

Acetaminophen - Pain reliever/Fever reducer,
Diphenhydramine HCl - Antihistamine/Cough suppressant
Phenylephrine HCl - Nasal decongestant

• Fast Relief of Cold & Flu symptoms
• Soothing Vicks Vapors

Tamper Evident:
Do not use if inner sealed packet is torn or broken.

MADE IN MEXICO
DIST. BY:
PROCTER & GAMBLE,
CINCINNATI, OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents

69 PACKETS



Delightful & Great
Tasting with Honey
Lemon Flavor.

ACTUAL SIZE





Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 • more than 4 doses in 24 hrs, which is the maximum daily amount for this product.
 • taken with other drugs containing acetaminophen.
 • 3 or more alcoholic drinks every day while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin redness • blisters • rash
 If a skin reaction occurs, stop use and seek medical help right away.

See throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 • liver disease • heart disease
 • high blood pressure • thyroid disease
 • diabetes

Drug Facts (continued)

Warnings
 • trouble urinating due to enlarged prostate gland
 • cough that occurs with too much phlegm (mucus)
 • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
 • a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if
 • you get nervous, dizzy or sleepless
 • pain, nasal congestion, or cough gets worse or lasts more than 7 days
 • fever gets worse or lasts more than 3 days
 • redness or swelling is present
 • new symptoms occur
 • cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 • take only as directed • only use the dose cap provided
 • do not exceed 4 doses per 24 hrs
 adults & children 12 yrs & over 30 mL every 4 hrs
 children 4 to under 12 yrs ask a doctor
 children under 4 yrs do not use

Drug Facts (continued)

Other information
 • each 30 mL contains sodium 81 mg
 • store at no greater than 25°C and do not refrigerate

Inactive ingredients alcohol, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions? 1-800-362-1683

**Drug Facts (continued)****Warnings****Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 44 mg
- store at no greater than 25°C

Inactive ingredients citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-800-362-1683

90262782

**Drug Facts (continued)**

• cough persists for more than 7 days, comes back or occurs with a fever, rash, or persistent headache
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)**Warnings****Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**When using this product**

- do not use more than directed.
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever

Tamper Evident:

Do not use if printed shrinkband seal around the neck is broken or missing
← **PEEL BACK FOR DRUG FACTS**

Drug Facts

Active ingredients (in each 15 mL) **Purpose**
Dextromethorphan HBr 10 mg.....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses temporarily relieves common cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202

Patents: www.pg.com/patents
www.vicks.com



P&G



www.pg.com

90402378









VICK'S

NyQuil[®]

VapoCOOL™

PARENTS:
Learn about non-medical abuse at
[www.StopMedicineAbuse.org](#)

ACTUAL SIZE

TAMPER EVIDENT:
This package is safety-sealed &
child resistant. Use only if blisters are intact.
If difficult to open, use scissors.

DIST. BY
PROCTER & GAMBLE,
CINCINNATI OH 45202

P&G
[www.pg.com](#)
Patents:
[www.pg.com/patents](#)
90778947

Eco-friendly packaging made from recycled materials.

DO NOT REUSE OR INCINERATE THIS PRODUCT AT ANY TIME.

©2011 P&G LLC

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Sleep aid/Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation • nasal congestion • sinus congestion & pressure
- headache • fever • sore throat • runny nose & sneezing • minimizes aches & pains
- temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 8 caplets in 24 hours, which is the maximum daily amount for this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness • hives • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ...with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product:

- do not use more than directed • excitability may occur, especially in children
- mild drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts

This could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.





